

Statistical Modeling: Propensity modeling

Cornell STSCI / INFO / ILRST 3900
causal3900.github.io

Oct 22, 2026

Learning goals for today

At the end of class, you will be able to

- ▶ Estimate average causal effects with a model for the treatment
- ▶ Reason about the bias variance tradeoff
- ▶ Use the Augmented IPW estimator to guard against model misspecification

After class:

- ▶ Hernán and Robins 2020 Chapter 12.1–12.2, Technical Points 13.2 and 13.3

G-formula: Outcome model

Standardization estimator

$$\hat{E}(Y^{a=1}) = \frac{1}{n} \sum_i \hat{E}(Y \mid L = \ell_i, A = 1)$$

G-formula: Outcome model

Standardization estimator

$$\hat{E}(Y^{a=1}) = \frac{1}{n} \sum_i \hat{E}(Y \mid L = \ell_i, A = 1)$$

Learn a model to predict Y given L and A

- ▶ Linear models potentially with interaction terms
- ▶ Other types of regression: logistic regression, poisson regression, etc
- ▶ Machine learning models

G-formula: Outcome model

Standardization estimator

$$\hat{E}(Y^{a=1}) = \frac{1}{n} \sum_i \hat{E}(Y \mid L = \ell_i, A = 1)$$

Learn a model to predict Y given L and A

- ▶ Linear models potentially with interaction terms
- ▶ Other types of regression: logistic regression, poisson regression, etc
- ▶ Machine learning models

For every unit i ,

- ▶ Set the treatment value to $A = 1$
- ▶ Predict the outcome

Then average over all units

Bias Variance trade-off

Bias and variance in making cakes:




Figure: High Bias, low variance



Figure: Low bias, High variance


Bias and variance in choosing causal model

- ▶ More complex models require a larger sample size to fit well
- ▶ Roughly speaking, more complex means that the model is more difficult to describe
- ▶ More complex can also mean we allow for non-linear relationships or also that we include more covariates




Bias and variance in choosing causal model

- ▶ More complex models require a larger sample size to fit well
- ▶ Roughly speaking, more complex means that the model is more difficult to describe
- ▶ More complex can also mean we allow for non-linear relationships or also that we include more covariates




Bias and variance in choosing causal model

- ▶ More complex models require a larger sample size to fit well
- ▶ Roughly speaking, more complex means that the model is more difficult to describe
- ▶ More complex can also mean we allow for non-linear relationships or also that we include more covariates



Bias and variance in choosing causal model

- ▶ More complex models require a larger sample size to fit well
- ▶ Roughly speaking, more complex means that the model is more difficult to describe
- ▶ More complex can also mean we allow for non-linear relationships or also that we include more covariates




Bias and variance in choosing causal model

- ▶ Is a a DAG ever “truly correct”?
- ▶ Can always add more confounders
- ▶ Would the bias from the confounders you could add substantially change your claim?
- ▶ Including additional confounders makes estimation more difficult, requires a larger sample to fit well

Inverse probability weighting


Inverse probability weighting



Observe:

In order to get the $ATE = E(Y^{a=1}) - E(Y^{a=0})$

Want: a world where everyone is treated



Want: a world where everyone is not treated



IPW: Everything together

General rule: Each individual should be reweighted by

$$\text{Weight} = \frac{1}{\text{Prob of their obs treatment status given their age}}$$

- ▶ Propensity Score: $\pi_i = \text{Pr}(A_i = 1 \mid L_i)$
- ▶ Let N be the number of total observations
- ▶ Hypothetical world where everyone is treated

$$E(Y^{a=1}) = \frac{1}{N} \sum_{i:A_i=1} \frac{1}{\pi_i} \times Y_i$$

- ▶ Hypothetical world where everyone is not treated

$$E(Y^{a=0}) = \frac{1}{N} \sum_{i:A_i=0} \frac{1}{1 - \pi_i} \times Y_i$$

$$\text{ATE} = \underbrace{\frac{1}{N} \sum_{i:A_i=1} \frac{1}{\pi_i} \times Y_i}_{\text{Weighted avg of treated obs}} - \underbrace{\frac{1}{N} \sum_{i:A_i=0} \frac{1}{1 - \pi_i} \times Y_i}_{\text{Weighted Avg of untreated obs}}$$

IPW: with data

Model the treatment assignment

$$\hat{\pi}_i = \hat{P}(A = 1 \mid L) = \text{logit}^{-1}(\hat{\alpha} + \hat{\gamma}L)$$

Estimate by inverse probability weighting

$$\hat{E}(Y^{a=1}) = \frac{1}{N} \sum_{i:A_i=1} \frac{Y_i}{\hat{\pi}_i}$$
$$\hat{E}(Y^{a=0}) = \frac{1}{N} \sum_{i:A_i=0} \frac{Y_i}{1 - \hat{\pi}_i}$$

Inverse probability of treatment weighting

$$\begin{aligned}\hat{E}(Y^{a=1}) &= \frac{1}{N} \sum_{i:A_i=1} \frac{Y_i}{\hat{\pi}_i} \\ &= \frac{1}{N} \sum_{i:A_i=1} \frac{\textcolor{red}{A}_i Y_i}{\hat{\pi}_i} && \text{(people with } A_i = 1\text{)} \\ &= \frac{1}{N} \left(\sum_{i:A_i=1} \frac{A_i Y_i}{\hat{\pi}_i} + \underbrace{\sum_{i:A_i=0} \frac{\textcolor{red}{A}_i Y_i}{\hat{\pi}_i}}_{=0} \right) && \text{(people with } A_i = 0\text{)} \\ &= \frac{1}{N} \sum_i \frac{A_i Y_i}{\hat{\pi}_i} && \text{(everyone)}\end{aligned}$$

(1)

Inverse probability of treatment weighting

$$\begin{aligned}\hat{E}(Y^{a=0}) &= \frac{1}{N} \sum_{i:A_i=0} \frac{Y_i}{1 - \hat{\pi}_i} \\&= \frac{1}{N} \sum_{i:A_i=0} \frac{(1 - A_i) Y_i}{\hat{\pi}_i} && \text{(people with } A_i = 0\text{)} \\&= \frac{1}{N} \left(\sum_{i:A_i=0} \frac{(1 - A_i) Y_i}{1 - \hat{\pi}_i} + \underbrace{\sum_{i:A_i=1} \frac{(1 - A_i) Y_i}{\hat{\pi}_i}}_{=0} \right) && \text{(people with } A_i = 1\text{)} \\&= \frac{1}{N} \sum_i \frac{(1 - A_i) Y_i}{1 - \hat{\pi}_i} && \text{(everyone)}\end{aligned}$$

(2)

Parametric model: propensity model

Model the treatment assignment

$$\hat{\pi}_i = \hat{P}(A = 1 \mid L) = \text{logit}^{-1}(\hat{\alpha} + \hat{\gamma}L)$$

Estimate by inverse probability weighting (IPW)

$$\hat{E}(Y^1) - \hat{E}(Y^0) = \frac{1}{N} \left(\sum_i \frac{A_i Y_i}{\hat{\pi}_i} - \sum_i \frac{(1 - A_i) Y_i}{1 - \hat{\pi}_i} \right)$$

Standardization vs Propensity score modeling

- ▶ When our predictions are exactly right, the two are equivalent
 - ▶ $\hat{E}(Y | A, L) = E(Y | A, L)$
 - ▶ $\hat{P}(A = a | L) = P(A = a | L)$
- ▶ Standardization (g-formula) is probably more popular, but it's not inherently better
- ▶ If outcomes easier to predict than treatment status, use standardization
- ▶ If treatment is easier to predict than outcomes, use IPW

Standardization vs Propensity score modeling

- ▶ When our predictions are exactly right, the two are equivalent
 - ▶ $\hat{E}(Y | A, L) = E(Y | A, L)$
 - ▶ $\hat{P}(A = a | L) = P(A = a | L)$
- ▶ Standardization (g-formula) is probably more popular, but it's not inherently better
- ▶ If outcomes easier to predict than treatment status, use standardization
- ▶ If treatment is easier to predict than outcomes, use IPW
- ▶ What if our statistical models are wrong?
 - ▶ The statistical model is not complex enough to accurately capture the true relationship?
 - ▶ What if we don't include the right covariates?

Augmented IPW

- ▶ Augmented IPW uses both estimates for outcome and propensity scores together
- ▶ Has a “double robustness” property

$$\hat{E}(Y^1) = \frac{1}{N} \left(\underbrace{\sum_i \hat{E}(Y \mid A = 1, L = \ell_i)}_{\text{g-computation}} + \underbrace{\sum_i \frac{A_i Y_i}{\hat{\pi}_i}}_{\text{IPW}} - \underbrace{\sum_i \frac{A_i}{\hat{\pi}_i} \hat{E}(Y \mid A = 1, L = \ell_i)}_{\text{new term}} \right)$$

Augmented IPW

Why is this a good idea?

- Standardization with the true conditional expectation would recover $E(Y^{a=1})$ when the sample size is large

$$\frac{1}{N} \sum_i E(Y \mid A = 1, L = \ell_i)$$

- AIPW estimates $E(Y^{a=1})$ by

$$\hat{E}(Y^{a=1}) = \frac{1}{N} \sum_i \left(\hat{E}(Y \mid A = 1, L = \ell_i) + \frac{A_i}{\hat{\pi}_i} Y_i - \frac{A_i}{\hat{\pi}_i} \hat{E}(Y \mid A = 1, L = \ell_i) \right)$$

Augmented IPW

Why is this a good idea?

- Standardization with the true conditional expectation would recover $E(Y^{a=1})$ when the sample size is large

$$\frac{1}{N} \sum_i E(Y \mid A = 1, L = \ell_i)$$

- AIPW estimates $E(Y^{a=1})$ by

$$\hat{E}(Y^{a=1}) = \frac{1}{N} \sum_i \left(\hat{E}(Y \mid A = 1, L = \ell_i) + \frac{A_i}{\hat{\pi}_i} Y_i - \frac{A_i}{\hat{\pi}_i} \hat{E}(Y \mid A = 1, L = \ell_i) \right)$$

- If the two terms inside are close to each other, then the AIPW will recover $E(Y^{a=1})$ when the sample size is large

Augmented IPW

- ▶ We want the expectation of the AIPW term to be close to the true conditional expectation

Augmented IPW

- ▶ We want the expectation of the AIPW term to be close to the true conditional expectation
- ▶ Note that $E(A_i) = Pr(A_i = 1 \mid L_i) = \pi_i$
- ▶ Fixing the predictions $\hat{E}(\cdot)$ and $\hat{\pi}$, take the expectation of AIPW term

$$\begin{aligned} E \left(\hat{E}(Y \mid A = 1, L = \ell_i) + \frac{A_i}{\hat{\pi}_i} Y_i - \frac{A_i}{\hat{\pi}_i} \hat{E}(Y \mid A = 1, L = \ell_i) \right) \\ = \hat{E}(Y \mid A = 1, L = \ell_i) + \frac{\pi_i}{\hat{\pi}_i} E(Y \mid A = 1, L = \ell_i) - \frac{\pi_i}{\hat{\pi}_i} \hat{E}(Y \mid A = 1, L = \ell_i) \end{aligned}$$

Augmented IPW

- ▶ We want the expectation of the AIPW term to be close to the true conditional expectation
- ▶ Note that $E(A_i) = Pr(A_i = 1 \mid L_i) = \pi_i$
- ▶ Fixing the predictions $\hat{E}(\cdot)$ and $\hat{\pi}$, take the expectation of AIPW term

$$\begin{aligned} E \left(\hat{E}(Y \mid A = 1, L = \ell_i) + \frac{A_i}{\hat{\pi}_i} Y_i - \frac{A_i}{\hat{\pi}_i} \hat{E}(Y \mid A = 1, L = \ell_i) \right) \\ = \hat{E}(Y \mid A = 1, L = \ell_i) + \frac{\pi_i}{\hat{\pi}_i} E(Y \mid A = 1, L = \ell_i) - \frac{\pi_i}{\hat{\pi}_i} \hat{E}(Y \mid A = 1, L = \ell_i) \\ = \hat{E}(Y \mid A = 1, L = \ell_i) + \frac{\pi_i}{\hat{\pi}_i} \left[E(Y \mid A = 1, L = \ell_i) - \hat{E}(Y \mid A = 1, L = \ell_i) \right] \end{aligned}$$

Augmented IPW

If the predicted outcome is correct:

$$\hat{E}(Y \mid A = 1, L = \ell_i) = E(Y \mid A = 1, L = \ell_i);$$

AIPW term

$$\begin{aligned} &= \hat{E}(Y \mid A = 1, L = \ell_i) + \frac{\pi_i}{\hat{\pi}_i} \left[E(Y \mid A = 1, L = \ell_i) - \hat{E}(Y \mid A = 1, L = \ell_i) \right] \\ &= E(Y \mid A = 1, L = \ell_i) + \frac{\pi_i}{\hat{\pi}_i} \underbrace{\left[E(Y \mid A = 1, L = \ell_i) - E(Y \mid A = 1, L = \ell_i) \right]}_{=0} \\ &= E(Y \mid A = 1, L = \ell_i) \end{aligned} \tag{3}$$

Augmented IPW

If the predicted outcome is correct:

$$\hat{E}(Y \mid A = 1, L = \ell_i) = E(Y \mid A = 1, L = \ell_i);$$

AIPW term

$$\begin{aligned} &= \hat{E}(Y \mid A = 1, L = \ell_i) + \frac{\pi_i}{\hat{\pi}_i} \left[E(Y \mid A = 1, L = \ell_i) - \hat{E}(Y \mid A = 1, L = \ell_i) \right] \\ &= E(Y \mid A = 1, L = \ell_i) + \frac{\pi_i}{\hat{\pi}_i} \underbrace{\left[E(Y \mid A = 1, L = \ell_i) - E(Y \mid A = 1, L = \ell_i) \right]}_{=0} \\ &= E(Y \mid A = 1, L = \ell_i) \end{aligned} \tag{3}$$

Regardless of how different $\hat{\pi}$ and π are!

Augmented IPW

If the predicted propensity score is correct $\hat{\pi} = \pi$:

AIPW term

$$\begin{aligned} &= \hat{E}(Y \mid A = 1, L = \ell_i) + \frac{\pi_i}{\hat{\pi}_i} \left[E(Y \mid A = 1, L = \ell_i) - \hat{E}(Y \mid A = 1, L = \ell_i) \right] \\ &= \hat{E}(Y \mid A = 1, L = \ell_i) + \underbrace{\frac{\pi_i}{\pi_i}}_{=1} \left[E(Y \mid A = 1, L = \ell_i) - \hat{E}(Y \mid A = 1, L = \ell_i) \right] \\ &= \hat{E}(Y \mid A = 1, L = \ell_i) + \left[E(Y \mid A = 1, L = \ell_i) - \hat{E}(Y \mid A = 1, L = \ell_i) \right] \\ &= E(Y \mid A = 1, L = \ell_i) \end{aligned} \tag{4}$$

Augmented IPW

If the predicted propensity score is correct $\hat{\pi} = \pi$:

AIPW term

$$\begin{aligned} &= \hat{E}(Y \mid A = 1, L = \ell_i) + \frac{\pi_i}{\hat{\pi}_i} \left[E(Y \mid A = 1, L = \ell_i) - \hat{E}(Y \mid A = 1, L = \ell_i) \right] \\ &= \hat{E}(Y \mid A = 1, L = \ell_i) + \underbrace{\frac{\pi_i}{\pi_i}}_{=1} \left[E(Y \mid A = 1, L = \ell_i) - \hat{E}(Y \mid A = 1, L = \ell_i) \right] \\ &= \hat{E}(Y \mid A = 1, L = \ell_i) + \left[E(Y \mid A = 1, L = \ell_i) - \hat{E}(Y \mid A = 1, L = \ell_i) \right] \\ &= E(Y \mid A = 1, L = \ell_i) \end{aligned} \tag{4}$$

Regardless of how different $\hat{E}(Y \mid A = 1, L = \ell_i)$ and $E(Y \mid A = 1, L = \ell_i)$ are!

Augmented IPW

- ▶ If you only use g-formula then outcome model must be correct
- ▶ If you only use IPW then propensity score model must be correct

Augmented IPW

- ▶ If you only use g-formula then outcome model must be correct
- ▶ If you only use IPW then propensity score model must be correct
- ▶ AIPW recovers $E(Y^1)$ if **either** the outcome model or the propensity score model is correct
- ▶ You get shots to be correct!
- ▶ One can be completely wrong as long as the other is correct

Augmented IPW

Double robustness allows a single “failure”



¹Daniel Craig in Knives Out

Augmented IPW

Augmented IPW uses both estimates for outcome and propensity scores together

$$\hat{E}(Y^{a=1}) = \frac{1}{N} \left(\underbrace{\sum_i \hat{E}(Y \mid A = 1, L = \ell_i)}_{\text{g-computation}} + \underbrace{\sum_i \frac{A_i Y_i}{\hat{\pi}_i}}_{\text{IPW}} - \underbrace{\sum_i \frac{A_i}{\hat{\pi}_i} \hat{E}(Y \mid A = 1, L = \ell_i)}_{\text{new term}} \right)$$

$$\hat{E}(Y^{a=0}) = \frac{1}{N} \left(\underbrace{\sum_i \hat{E}(Y \mid A = 0, L = \ell_i)}_{\text{g-computation}} + \underbrace{\sum_i \frac{(1 - A_i) Y_i}{1 - \hat{\pi}_i}}_{\text{IPW}} - \underbrace{\sum_i \frac{1 - A_i}{1 - \hat{\pi}_i} \hat{E}(Y \mid A = 0, L = \ell_i)}_{\text{new term}} \right)$$

Augmented IPW

Augmented IPW uses both estimates for outcome and propensity scores together

$$\hat{E}(Y^{a=1}) = \frac{1}{N} \left(\underbrace{\sum_i \hat{E}(Y \mid A = 1, L = \ell_i)}_{\text{g-computation}} + \underbrace{\sum_i \frac{A_i Y_i}{\hat{\pi}_i}}_{\text{IPW}} - \underbrace{\sum_i \frac{A_i}{\hat{\pi}_i} \hat{E}(Y \mid A = 1, L = \ell_i)}_{\text{new term}} \right)$$

$$\hat{E}(Y^{a=0}) = \frac{1}{N} \left(\underbrace{\sum_i \hat{E}(Y \mid A = 0, L = \ell_i)}_{\text{g-computation}} + \underbrace{\sum_i \frac{(1 - A_i) Y_i}{1 - \hat{\pi}_i}}_{\text{IPW}} - \underbrace{\sum_i \frac{1 - A_i}{1 - \hat{\pi}_i} \hat{E}(Y \mid A = 0, L = \ell_i)}_{\text{new term}} \right)$$

Estimate ATE

$$\hat{E}(Y^{a=1}) - \hat{E}(Y^{a=0})$$

Augmented IPW

- ▶ Double robustness: robust against misspecification of either (but not both)
- ▶ If the outcome model is well specified, using standardization with just the outcome model often has smaller variance
- ▶ If the outcome model is not well specified, using standardization with just the outcome model may not be correct even as the sample size grows very large
- ▶ Using AIPW provides insurance against misspecification

Learning goals for today

At the end of class, you will be able to

- ▶ Estimate average causal effects with a model for the treatment
- ▶ Reason about the bias variance tradeoff
- ▶ Use the Augmented IPW estimator to guard against model misspecification

After class:

- ▶ Hernán and Robins 2020 Chapter 12.1–12.2, Technical Points 13.2 and 13.3